

On special values of spinor L -functions of Siegel cusp eigenforms of genus 3

Francesco Chiera Kirill Vankov

May 14, 2008

Abstract

We compute the special values for the spinor L -function $L(s, F_{12}, \text{spin})$ in the critical strip $s = 12, \dots, 19$, where F_{12} is the unique (up to a scalar) Siegel cusp form of degree 3 and weight 12, which was constructed by Miyawaki. These values are proportional to the product of Petersson inner products of Ramanujan's Δ by itself and the cusp form of weight 20 for $\text{SL}_2(\mathbb{Z})$ by itself by a rational number and some power of π . We also verify this result numerically using Dokchitser's ComputeL PARI package. To our knowledge this is the first example of a spinor L -function of Siegel cusp forms of degree 3, when the special values can be computed explicitly.

Contents

1	Introduction	2
2	Generalities and notations	3
3	The expression for $L(s, \Delta) L(s-1, \Delta)$	5
4	Computation of $L(s, \Delta \otimes G_{2,2})$	7
5	Result for $L(s-9, \Delta) L(s-10, \Delta)$	12
6	Computation of $L(s, \Delta \otimes g_{20})$	13
7	The main identity	15
8	Numerical computation of Petersson inner product	16
9	Numerical verification	17

1 Introduction

In the remarkable paper [Miy92] I. Miyawaki considered certain Siegel cusp forms of degree 3, and on the basis of some numerical calculations, he was able to make interesting conjectures about the degeneration of the standard and spinor L -functions associated to such cusp forms. Several years later T. Ikeda [Ike06] proved Miyawaki's conjecture related to the standard L -function. Basically he was able to construct an explicit lifting from Siegel cusp forms of degree r to Siegel cusp forms of degree $r + 2n$. In particular, it turns out that the only cusp form of degree 3 and weight 12 is a basic example of this lifting (for $r = 1, n = 1$).

Recall that Miyawaki constructed his numerical examples by means of theta functions with spherical functions. Namely, let E_8 be the unique even unimodular lattice of rank 8 i.e.,

$$E_8 = \left\{ {}^t(x_1, \dots, x_8) \in \mathbb{R}^8 \left| \begin{array}{l} 2x_i \in \mathbb{Z} (i = 1, \dots, 8), \\ x_1 + \dots + x_8 \in 2\mathbb{Z}, \\ x_i - x_j \in \mathbb{Z} \end{array} \right. \right\}, \quad (1)$$

and

$$Q = \begin{pmatrix} 1 & 0 & 0 & i & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & i & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & i & 0 & 0 \end{pmatrix} \quad (2)$$

be 3×8 matrix. Then the theta series

$$\begin{aligned} F_{12} &= \sum_{v_1, v_2, v_3 \in E_8} \Re(\det(Q \cdot (v_1, v_2, v_3))^8 \exp(\pi i \sigma(\langle v_i, v_j \rangle Z))) \\ &= \sum_{N > 0} a(N) \exp(2\pi i \sigma(NZ)) \end{aligned} \quad (3)$$

is a cusp form of weight 12 with respect to $\mathrm{Sp}_3(\mathbb{Z})$, where

$$\mathrm{Sp}_n(\mathbb{Z}) = \left\{ M \in \mathrm{Mat}_{2n} \mid M \begin{pmatrix} 0 & \mathrm{I}_n \\ -\mathrm{I}_n & 0 \end{pmatrix} {}^t M = \begin{pmatrix} 0 & \mathrm{I}_n \\ -\mathrm{I}_n & 0 \end{pmatrix} \right\}. \quad (4)$$

In the recent work [Hei08] B. Heim proved Miyawaki's conjecture relevant to the spinor L -function for the specific case of the cusp form F_{12} . In fact, Heim has showed that the following equality holds:

$$L(s, F_{12}, \text{spin}) = L(s - 9, \Delta) L(s - 10, \Delta) L(s, \Delta \otimes g_{20}), \quad (5)$$

where Δ is Ramanujan's discriminant cusp form and g_{20} is the cusp form of weight 20 of level 1. This theorem is the starting point of our investigation. The purpose of this note is to show that one can provide the explicit rational number R_s and power of π such that for each critical value $s = 12, \dots, 19$

$$L(s, F_{12}, \text{spin}) = R_s \pi^{\alpha_s} \langle \Delta, \Delta \rangle \langle g_{20}, g_{20} \rangle. \quad (6)$$

We also compute the values numerically using the SAGE software [SAG] and Dokchitser's COMPUTEL PARI package [Dok, Dok04].

To our knowledge, this is the first example of a spinor L -function of Siegel cusp forms of degree 3, when the special values can be computed explicitly.

It is possible to apply the same technique to compute the critical values of the spinor L -functions for non cuspidal modular forms; but there exist direct (and more simple) methods in this case. For example, due to Zharkovskaya [Žar74, Theorem 1] there is the famous equality for non cuspidal forms

$$L(s, F, \text{spin}) = L(s, \Phi(F), \text{spin}) L(s - k + n, \Phi(F), \text{spin}), \quad (7)$$

where Φ is the Siegel operator. We could also use our approach to compute the conjectural special values of the spinor L -function $L(s, F_{14}, \text{spin})$ for the unique Siegel cusp for of degree 3 and weight 14.

The authors are very grateful to Alexei Panchishkin for intensive and encouraging discussions, this work was undertaken due to his keen interest into the subject. We also express our thanks to Bernhard Heim for useful remarks.

2 Generalities and notations

Let $\mathfrak{H} = \{z \in \mathbb{C} \mid \Im(z) > 0\}$ be the upper half-plane. For a positive integer k and a Dirichlet character χ modulo a positive integer N such that $\chi(-1) = (-1)^k$, we denote by $\mathcal{M}_k(\Gamma_0(N), \chi)$ the vector space of all holomorphic modular forms $f(z)$ of weight k satisfying

$$f(\gamma(z)) = \chi(d)(cz + d)^k f(z) \text{ for all } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N), \quad (8)$$

where the variable $z \in \mathfrak{H}$, $\gamma(z) = \frac{az + b}{cz + d}$, and

$$\Gamma_0(N) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}_2(\mathbb{Z}) \mid c \equiv 0 \pmod{N} \right\}. \quad (9)$$

We denote by $S_k(N, \chi)$ the subspace of $\mathcal{M}_k(\Gamma_0(N), \chi)$ consisting of all cusp forms. Every element f of $\mathcal{M}_k(\Gamma_0(N), \chi)$ has a Fourier expansion

$$f(z) = \sum_{n=0}^{\infty} a(n) q^n, \quad (10)$$

where $q = \exp(2\pi iz)$ and $a(n)$ are complex numbers in general.

The L -function associated to f is defined as $L(s, f) = \sum_{n=1}^{\infty} a(n) n^{-s}$. More generally, with an arbitrary Dirichlet character ω , the twisted L -function is defined as $L(s, f, \omega) = \sum_{n=1}^{\infty} a(n) \omega(n) n^{-s}$. These L -functions can be also written in the form of Euler product:

$$L(s, f) = \prod_{p \text{ prime}} (1 - a(p) p^{-s} + \chi(p) p^{k-1-2s})^{-1}, \quad (11)$$

$$L(s, f, \omega) = \prod_{p \text{ prime}} (1 - a(p) \omega(p) p^{-s} + \chi(p) \omega(p)^2 p^{k-1-2s})^{-1}. \quad (12)$$

Next, the Dirichlet L -series for any character χ of conductor N is given by

$$L(s, \chi) = \sum_{n=1}^{\infty} \chi(n) n^{-s} \quad (13)$$

and its Euler product

$$L(s, \chi) = \prod_{p \nmid N} \frac{1}{1 - \chi(p) p^{-s}}. \quad (14)$$

In the case, when χ is the identity Dirichlet character, the latter series is Riemann's zeta function $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$.

Let $g(z) = \sum_{n=0}^{\infty} b(n) q^n \in \mathcal{M}_l(\Gamma_0(N), \xi)$ be another modular form of weight l with Fourier coefficients $b(n)$. The L function associated to two modular forms f and g is given by the additive convolution

$$L(s, f, g) = \sum_{n=1}^{\infty} a(n) b(n) n^{-s}. \quad (15)$$

Another type of L -functions associated to two modular forms is Rankin's product L -function (multiplicative convolution). It is denoted by $L(s, f \otimes g)$ and defined as (see [Shi76, page 786]):

$$L(s, f \otimes g) = L_N(2s + 2 - k - l, \chi\xi) L(s, f, g), \quad (16)$$

where $L_N(s, \omega)$ with a Dirichlet character ω modulo N is defined, as usual, by (13) with $\omega(n) = 0$ for $(n, N) \neq 1$, and the Euler factors in (14), corresponding to the prime divisors of a number N , have been omitted. Note, that in the case, when f and g are cusp eigenforms, the lefthand side of (16) is an Euler product of degree 4 in view of the following Lemma:

LEMMA 2.1 (LEMMA 1, [Shi76]) *Suppose we have formally*

$$\sum_{n=1}^{\infty} A(n) n^{-s} = \prod_p [(1 - \alpha_p p^{-s})(1 - \alpha'_p p^{-s})]^{-1}$$

and

$$\sum_{n=1}^{\infty} B(n) n^{-s} = \prod_p [(1 - \beta_p p^{-s})(1 - \beta'_p p^{-s})]^{-1}.$$

Then

$$\sum_{n=1}^{\infty} A(n) B(n) n^{-s} = \frac{1 - \alpha\alpha'\beta\beta'p^{-2s}}{(1 - \alpha\beta p^{-s})(1 - \alpha\beta'p^{-s})(1 - \alpha'\beta p^{-s})(1 - \alpha'\beta'p^{-s})}.$$

For two elements $f, h \in \mathcal{M}_k(\Gamma_0(N))$ such that fh is a cusp form, the Petersson inner product $\langle f, h \rangle$ is defined as

$$\langle f, h \rangle = \frac{1}{[\mathrm{SL}_2(\mathbb{Z}) : \Gamma_0(N)]} \int_{\Phi_N} \overline{f(z)} h(z) y^{k-2} dx dy, \quad (17)$$

where $z = x + iy$, Φ_N is a fundamental domain for $\mathfrak{H}$ modulo $\Gamma_0(N)$ and the bar denotes the complex conjugate. We also define $\langle f, h \rangle$ by (17) for nearly holomorphic modular forms f and h on $\mathfrak{H}$ whenever the integral is convergent (see [Shi07, section 8.2] for definition and properties of the nearly holomorphic modular forms).

We also give a brief definition of the spinor L -function. For a Hecke eigenform F with local Satake parameters $\mu_0, \mu_1, \dots, \mu_n$ it is given by the infinite product

$$L(s, F, \text{spin}) = \prod_p L_p(s, F, \text{spin}), \quad (18)$$

where for each prime number p

$$L_p(s, F, \text{spin}) = \left((1 - \mu_0 p^{-s}) \prod_{r=1}^n \prod_{i_1 < \dots < i_r} (1 - \mu_0 \mu_{i_1} \dots \mu_{i_r} p^{-s}) \right)^{-1}. \quad (19)$$

We do not use this definition in our computation, but develop the right-hand side of the identity (5). The explicit description of the Satake parameters for F_{12} is given in [Hei08].

3 The expression for $L(s, \Delta) L(s-1, \Delta)$

Let $f = \Delta$ be Ramanujan's discriminant modular form of weight $k = 12$:

$$\begin{aligned} f(z) = \Delta(z) &= \sum_{n=1}^{\infty} \tau(n) q^n = q \prod_{n=1}^{\infty} (1 - q^n)^{24} \\ &= q - 24 q^2 + 252 q^3 - 1472 q^4 + 4830 q^5 - 6048 q^6 + \dots, \end{aligned} \quad (20)$$

where $\tau(n)$ is Ramanujan's tau function. The associated L -function is

$$L(s, f) = \sum_{n=1}^{\infty} \tau(n) n^{-s} = \prod_p (1 - \tau(p) p^{-s} + p^{11-2s})^{-1}. \quad (21)$$

Let $G_2(z) = -\frac{1}{24} + \sum_{n=1}^{\infty} \sigma_1(n) q^n = -\frac{1}{24} + q + 3q^2 + 4q^3 + 7q^4 + \dots$, where

$\sigma_1(n) = \sum_{d|n} d$ is the divisor function defined as the sum of the divisors

of n . Consider the Eisenstein series (see [Miy06, Lemma 7.2.19, (2)])

$$g(z) = G_{2,p}(z) = G_2(z) - p G_2(pz) = \frac{p-1}{24} + \sum_{n=1}^{\infty} \sum_{\substack{d|n \\ p \nmid d}} d q^n, \quad (22)$$

of weight 2 for $\Gamma_0(p)$ and the corresponding L series

$$\begin{aligned} L(s, g) &= \sum_{n=1}^{\infty} \sum_{\substack{d|n \\ p \nmid d}} d n^{-s} = \sum_{\substack{d, d_1 \geq 1 \\ p \nmid d}} d (d d_1)^{-s} = \sum_{\substack{d \geq 1 \\ p \nmid d}} d^{1-s} \sum_{d_1 \geq 1} d_1^{-s} \\ &= (1 - p^{1-s}) \zeta(s-1) \zeta(s). \end{aligned} \quad (23)$$

Let us put $p = 2$, then

$$\begin{aligned} g(z) &= G_{2,2}(z) = G_2(z) - 2G_2(2z) \\ &= \frac{1}{24} + q + q^2 + 4q^3 + q^4 + 6q^5 + 4q^6 + \cdots, \end{aligned} \quad (24)$$

and

$$L(s, g) = (1 - 2^{1-s}) \zeta(s-1) \zeta(s). \quad (25)$$

For $f = \Delta = \sum \tau(n) q^n \in S_{12}(2)$ and $g = G_{2,2} = \sum b(n) q^n \in \mathcal{M}_2(\Gamma_0(2), \xi)$ we have $k = 12$ and $l = 2$. Put $N = 2$, $\chi = 1$ and

$$\xi(n) = (1 \bmod N)(n) = \begin{cases} 1, & \text{if } n \text{ odd;} \\ 0, & \text{if } n \text{ even.} \end{cases} \quad (26)$$

Assume $1 - \tau(p)X + p^{11}X^2 = (1 - \alpha_p X)(1 - \alpha'_p X)$, then $\alpha_p + \alpha'_p = \tau(p)$, $\alpha_p \alpha'_p = p^{11}$, and

$$L(s, f) = \prod_p ((1 - \alpha_p p^{-s})(1 - \alpha'_p p^{-s}))^{-1}. \quad (27)$$

Similarly, consider

$$\begin{aligned} L(s, g) &= \sum_{n=1}^{\infty} b(n) n^{-s} \\ &= (1 - 2^{1-s}) \zeta(s-1) \zeta(s) \\ &= (1 - 2^{1-s}) \prod_p ((1 - p^{1-s})(1 - p^{-s}))^{-1} \\ &= \prod_p ((1 - \beta_p p^{-s})(1 - \beta'_p p^{-s}))^{-1}, \end{aligned} \quad (28)$$

where $\beta(p) = 1$ for all p , $\beta'(2) = 0$ and $\beta'(p) = p$ for all odd primes. By definition and using Lemma 2.1

$$\begin{aligned}
L(s, \Delta \otimes G_{2,2}) &= L_2(2s+2-12-2, \psi) L(s, f, g) \\
&= \prod_{p \neq 2} (1 - p^{12-2s})^{-1} \cdot \sum_{n=1}^{\infty} \tau(n) b(n) n^{-s} \\
&= \prod_{p \neq 2} (1 - p^{12-2s})^{-1} \times \\
&\quad \times \prod_p \frac{1 - \alpha_p \alpha'_p \beta_p \beta'_p p^{-2s}}{(1 - \alpha_p \beta_p p^{-s}) (1 - \alpha'_p \beta_p p^{-s}) (1 - \alpha_p \beta'_p p^{-s}) (1 - \alpha'_p \beta'_p p^{-s})} \\
&= \frac{1}{(1 - \alpha_2 2^{-s}) (1 - \alpha'_2 2^{-s})} \times \\
&\quad \times \prod_{p \neq 2} \frac{(1 - p^{12-2s})^{-1} (1 - p^{11} p p^{-2s})}{(1 - \alpha_p p^{-s}) (1 - \alpha'_p p^{-s}) (1 - \alpha_p p p^{-s}) (1 - \alpha'_p p p^{-s})} \\
&= \frac{1}{(1 - \alpha_2 2^{-s}) (1 - \alpha'_2 2^{-s})} \times \\
&\quad \times \prod_{p \neq 2} \frac{1}{((1 - \alpha_p p^{-s}) (1 - \alpha'_p p^{-s})) ((1 - \alpha_p p^{1-s}) (1 - \alpha'_p p^{1-s}))} \\
&= \prod_p \frac{1}{(1 - \alpha_p p^{-s}) (1 - \alpha'_p p^{-s})} \prod_{p \neq 2} \frac{1}{(1 - \alpha_p p^{1-s}) (1 - \alpha'_p p^{1-s})} \\
&= L(s, \Delta) L(s-1, \Delta) (1 - \tau(2) 2^{1-s} + 2^{11} s^{2-2s}) \\
&= (1 + 3 \cdot 2^{4-s} + s^{13-2s}) L(s, \Delta) L(s-1, \Delta).
\end{aligned} \tag{29}$$

Finally, we obtain the following identity

$$L(s, \Delta) L(s-1, \Delta) = \frac{L(s, \Delta \otimes G_{2,2})}{(1 + 3 \cdot 2^{4-s} + 2^{13-2s})}. \tag{30}$$

4 Computation of $L(s, \Delta \otimes G_{2,2})$

Now we express $L(s, \Delta \otimes G_{2,2})$ (at integral points $s = 3, \dots, 10$) as a multiple of Petersson inner product $\langle \Delta, \Delta \rangle$. Using [Shi76, (2.4)],

$$\begin{aligned}
L(s, \Delta \otimes G_{2,2}) &= \frac{(4\pi)^s}{2\Gamma(s)} \int_{\Phi_2} \overline{\Delta(z)} G_{2,2}(z) E_{10,2}(z, s-11, \xi) y^{s-1} dx dy \\
&= \frac{(4\pi)^s}{2\Gamma(s)} \int_{\Phi_2} \overline{\Delta(z)} G_{2,2}(z) E_{10,2}(z, s-11, \xi) y^{s-11} y^{10} dx dy \\
&= \frac{(4\pi)^s [\text{SL}_2(\mathbb{Z}) : \Gamma_0(2)]}{2\Gamma(s)} \langle \Delta(z), G_{2,2}(z) y^{s-11} E_{10,2}(z, s-11, \xi) \rangle \tag{31} \\
&= \frac{3}{2} \frac{(4\pi)^s}{\Gamma(s)} \langle \Delta(z), \mathcal{H}ol(G_{2,2}(z) y^{s-11} E_{10,2}(z, s-11, \xi)) \rangle \\
&= \frac{3}{2} \frac{(4\pi)^{11}}{\Gamma(s)} \langle \Delta(z), \mathcal{H}ol(G_{2,2}(z) (4\pi y)^{s-11} E_{10,2}(z, s-11, \xi)) \rangle,
\end{aligned}$$

where $\langle f, g \rangle$ is the Petersson inner product (17), Φ_2 denotes a fundamental domain for $\Gamma_0(2) \backslash \mathfrak{H}$, $z = x + iy$,

$$E_{\lambda, N}(z, s, \xi) = \sum'_{(m, n)} \xi(n) (mNz + n)^{-\lambda} |mNz + n|^{-2s}, \quad (32)$$

$\sum'$ denotes the summation over all $(m, n) \in \mathbb{Z}^2$, $(m, n) \neq (0, 0)$, $\mathcal{H}ol(F)$ is the operator of holomorphic projection (see [CP04, (2.148)]). It is defined so, that $\langle f, F \rangle = \langle f, \mathcal{H}ol(F) \rangle$ for all $f \in S_k(N, \psi)$.

In order to compute the holomorphic projection of the product in the last identity of (31) $\mathcal{H}ol(G_{2,2}(z) (4\pi y)^{s-11} E_{10,2}(z, s-11, \xi))$ we write the Fourier expansion for $E_{10,2}(z, s-11, \xi)$ in the convenient form using the Whittaker functions by applying the Proposition 2.2 of [Pan03], where in notations of that proposition

$$s \in \{3, \dots, 10\}, \quad s-11 < 0, \quad s-11+10 = s-1 > 0,$$

$$a = 0, \quad b = 1,$$

$$\mathbf{E}_{10,2}(z, s-11, 0, 1) = E_{10,2}(z, s-11, \xi),$$

$$\delta\left(\frac{a}{N}\right) = 1,$$

$$\zeta(s; 0, 2) = \sum_{0 < n \equiv 0(2)} n^{-s} = \sum_{n=1}^{\infty} (2n)^{-s} = 2^{-s} \zeta(s),$$

$$\zeta(s; 1, 2) = \sum_{0 < n \equiv 1(2)} n^{-s} = \zeta(s) - \zeta(s; 0, 2) = (1 - 2^{-s}) \zeta(s).$$

The Whittaker function $W(y, \alpha, \beta)$ is defined as (see [Pan03, (2.4)], for example)

$$W(y, \alpha, \beta) = \Gamma(\beta)^{-1} \int_0^{+\infty} (u+1)^{\alpha-1} u^{\beta-1} e^{-yu} du \quad (33)$$

for $y > 0$, $\alpha, \beta \in \mathbb{C}$ with $\Re(\beta) > 0$ and for arbitrary α and β this function is defined by the analytic continuation and the functional equation:

$$W(y, \alpha, \beta) = y^{1-\alpha-\beta} W(y, 1-\beta, 1-\alpha). \quad (34)$$

For a non negative integer r , we have

$$W(y, \alpha, -r) = \sum_{i=0}^r \frac{(-1)^i \binom{r}{i} \Gamma(\alpha)}{\Gamma(\alpha-i)} y^{r-i}. \quad (35)$$

Therefore,

$$\begin{aligned}
& (4\pi y)^{s-11} E_{10,2}(z, s-11, \xi) \\
&= (4\pi y)^{s-11} 2 \zeta(2s-12; 1, 2) \\
&+ (4\pi y)^{s-11} \frac{(-2\pi i)^{2s-12} (-1)^{s-11} \Gamma(2s-13)}{(4\pi y)^{2s-13} 2 \Gamma(s-1) \Gamma(s-11)} 2 \zeta(2s-13; 0, 2) \\
&+ (4\pi y)^{s-11} \frac{(-2\pi i)^{2s-12} (-1)^{s-11}}{2^{2s-12} \Gamma(s-1)} \times \\
&\quad \times \sum_{n=1}^{\infty} \sum_{\pm d|n} \operatorname{sgn}(d) d^{2s-13} e^{\pi i d} W(4\pi n y, s-1, s-11) q^n \\
&= 2 (4\pi y)^{s-11} (1 - 2^{12-2s}) \zeta(2s-12) \\
&- (4\pi y)^{2-s} \frac{2\pi^{2s-12} \Gamma(2s-13) \zeta(2s-13)}{\Gamma(s-11) \Gamma(s-1)} \\
&- (4\pi y)^{s-11} \frac{2\pi^{2s-12}}{\Gamma(s-1)} \sum_{n=1}^{\infty} \sum_{d|n} (-1)^d d^{2s-13} W(4\pi n y, s-1, s-11) q^n.
\end{aligned} \tag{36}$$

Let

$$\begin{aligned}
C'_0 &= C'_0(s) = (-1) \frac{2\pi^{2s-12} \Gamma(2s-13) \zeta(2s-13)}{\Gamma(s-11) \Gamma(s-1)}, \\
C''_0 &= C''_0(s) = (2 - 2^{13-2s}) \zeta(2s-12), \\
C_1 &= C_1(s) = 2\pi^{2s-12}, \\
C_2 &= C_2(s) = (2 - 2^{2s-12}) \pi^{2s-12},
\end{aligned} \tag{37}$$

then

$$\begin{aligned}
& (4\pi y)^{s-11} E_{10,2}(z, s-11, \xi) \\
&= C'_0 (4\pi y)^{2-s} + C''_0 (4\pi y)^{s-11} \\
&\quad + C_1 \frac{W(4\pi y, s-1, s-11)}{\Gamma(s-1)} (4\pi y)^{s-11} q \\
&\quad + C_2 \frac{W(8\pi y, s-1, s-11)}{\Gamma(s-1)} (4\pi y)^{s-11} q^2 + \dots
\end{aligned} \tag{38}$$

Recall that $G_{2,2}(z) = 1/24 + q + q^2 + \dots$ by (24). We write the Fourier coefficients $\tilde{A}_i(s, y)$ for the product $F = G_{2,2}(z) (4\pi y)^{s-11} E_{10,2}(z, s-11, \xi) = \sum \tilde{A}_n(s, y) q^n$ in order to apply the Holomorphic Projection Lemma [GZ86, Proposition (5.1)] to find the image of the projection operator $\mathcal{H}ol(F) = \sum A_n(s) q^n$ (the Lemma is originally due to Sturm [Stu80]). It should be noted, that the relevant polynomial decay hypotheses of the Lemma are satisfied for all actions $E_{10,2}(z, s-11, \xi)|\gamma$ of $\gamma \in \operatorname{SL}_2(\mathbb{Z})$ and each critical point s , see [Pan03, (2.3)].

We need to compute just two coefficients $A_1 = A_1(s)$ and $A_2 = A_2(s)$ since the result of the holomorphic projection belongs to the space of cusp forms for the subgroup $\Gamma_0(2)$, which has the dimension 2. Then we find the linear combination representing $\mathcal{H}ol(F)$ in the basis $\{\Delta(z), \Delta(2z)\}$:

$$\mathcal{H}ol(F) = \alpha \cdot \Delta(z) + \beta \cdot \Delta(2z). \tag{39}$$

Namely, the computation of A_1 and A_2 gives:

$$\begin{aligned}
A_1 &= \frac{C'_0}{10!} \int_0^{+\infty} (4\pi y)^{2-s} e^{-4\pi y} (4\pi y)^{10} d(4\pi y) \\
&+ \frac{C''_0}{10!} \int_0^{+\infty} (4\pi y)^{s-11} e^{-4\pi y} (4\pi y)^{10} d(4\pi y) \\
&+ \frac{C_1}{24 \cdot 10!} \int_0^{+\infty} \frac{W(4\pi y, s-1, s-11)}{\Gamma(s-1)} (4\pi y)^{s-11} e^{-4\pi y} (4\pi y)^{10} d(4\pi y) \\
&= \frac{\Gamma(13-s)}{10!} C'_0 + \frac{\Gamma(s)}{10!} C''_0 \\
&+ \frac{C_1}{24 \cdot 10!} \int_0^{+\infty} \sum_{i=0}^{11-s} \frac{(-1)^i \binom{11-s}{i} (4\pi y)^{10-i}}{\Gamma(s-1-i)} e^{-4\pi y} d(4\pi y) \\
&= \frac{\Gamma(13-s)}{10!} C'_0 + \frac{\Gamma(s)}{10!} C''_0 \\
&+ \frac{C_1}{24 \cdot 10!} \sum_{i=0}^{11-s} \frac{(-1)^i \binom{11-s}{i} \Gamma(11-i)}{\Gamma(s-1-i)},
\end{aligned} \tag{40}$$

$$\begin{aligned}
A_2 &= \frac{C'_0}{10!} \int_0^{+\infty} (4\pi y)^{2-s} e^{-8\pi y} (8\pi y)^{10} d(8\pi y) \\
&+ \frac{C''_0}{10!} \int_0^{+\infty} (4\pi y)^{s-11} e^{-8\pi y} (8\pi y)^{10} d(8\pi y) \\
&+ \frac{C_1}{10!} \int_0^{+\infty} \frac{W(4\pi y, s-1, s-11)}{\Gamma(s-1)} (4\pi y)^{s-11} e^{-8\pi y} (8\pi y)^{10} d(8\pi y) \\
&+ \frac{C_2}{24 \cdot 10!} \int_0^{+\infty} \frac{W(8\pi y, s-1, s-11)}{\Gamma(s-1)} (4\pi y)^{s-11} e^{-8\pi y} (8\pi y)^{10} d(8\pi y) \\
&= \frac{C'_0}{10!} 2^{s-2} \int_0^{+\infty} (8\pi y)^{12-s} e^{-8\pi y} d(8\pi y) \\
&+ \frac{C''_0}{10!} 2^{11-s} \int_0^{+\infty} (8\pi y)^{s-1} e^{-8\pi y} d(8\pi y) \\
&+ \frac{C_1}{10!} \int_0^{+\infty} \sum_{i=0}^{11-s} \frac{(-1)^i \binom{11-s}{i} (4\pi y)^{11-s-i}}{\Gamma(s-1-i)} (4\pi y)^{s-11} e^{-8\pi y} (8\pi y)^{10} d(8\pi y) \\
&+ \frac{C_2}{24 \cdot 10!} \int_0^{+\infty} \sum_{i=0}^{11-s} \frac{(-1)^i \binom{11-s}{i} (8\pi y)^{11-s-i}}{\Gamma(s-1-i)} (4\pi y)^{s-11} e^{-8\pi y} (8\pi y)^{10} d(8\pi y) \\
&= \frac{\Gamma(13-s)}{10!} 2^{s-2} C'_0 + \frac{\Gamma(s)}{10!} 2^{11-s} C''_0 \\
&+ \frac{C_1}{10!} \sum_{i=0}^{11-s} 2^i \frac{(-1)^i \binom{11-s}{i} \Gamma(11-i)}{\Gamma(s-1-i)} \\
&+ \frac{C_2}{24 \cdot 10!} 2^{11-s} \sum_{i=0}^{11-s} \frac{(-1)^i \binom{11-s}{i} \Gamma(11-i)}{\Gamma(s-1-i)}.
\end{aligned} \tag{41}$$

These Fourier coefficients are rational numbers up to the factor π^{2s-12}

Table 1: Fourier coefficient of $\mathcal{Hol}(G_{2,2}(z)(4\pi y)^{s-11}E_{10,2}(z, s-11, \xi))$

s	π -factor	$A_1(s)$	$A_2(s)$
3	π^{-6}	$\frac{1}{50}$	$\frac{76}{25}$
4	π^{-4}	$\frac{1}{270}$	$\frac{56}{135}$
5	π^{-2}	$\frac{1}{1440}$	$\frac{1}{20}$
6	π^0	$\frac{1}{6048}$	$\frac{5}{756}$
7	π^2	$\frac{1}{16800}$	$\frac{1}{900}$
8	π^4	$\frac{17}{518400}$	$\frac{13}{64800}$
9	π^6	$\frac{11}{453600}$	$\frac{-1}{56700}$
10	π^8	$\frac{13}{604800}$	$\frac{-1}{10800}$

for each $s \in \{3, \dots, 10\}$, see Table 1.

Next, we compute α and β in the linear combination (39) by comparing the coefficients A_1 and A_2 corresponding to terms q and q^2 with the equivalent linear combination of coefficients of our basis functions $\Delta(z) = q - 24q^2 + \dots$ and $\Delta(2z) = q^2 + \dots$:

$$\begin{cases} A_1 = \alpha \cdot 1 + \beta \cdot 0 \\ A_2 = \alpha \cdot (-24) + \beta \cdot 1 \end{cases} \quad (42)$$

Resolving this system of linear equations, we obtain

$$\begin{aligned} \alpha &= A_1 \\ \beta &= 24 A_1 + A_2, \end{aligned} \quad (43)$$

therefore, we obtain the following identity for the Rankin's convolution of Δ and $G_{2,2}$ (31):

$$\begin{aligned} L(s, \Delta \otimes G_{2,2}) &= \frac{3}{2} \frac{(4\pi)^{11}}{\Gamma(s)} (A_1(s) \langle \Delta(z), \Delta(z) \rangle + (24 A_1(s) + A_2(s)) \langle \Delta(z), \Delta(2z) \rangle). \end{aligned} \quad (44)$$

We simplify the obtained expression even further. Recall that

$$\Delta(2z) = 2^{-k/2} \Delta(z) |V(2), \quad (45)$$

where $k = 12$ is the weight of Δ and $V(2)$ is the operator $\begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$. Then

$$\langle \Delta(z), \Delta(2z) \rangle = 2^{-6} \langle \Delta(z), \Delta(z) | \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} \rangle. \quad (46)$$

Consider $\gamma \in \Gamma_0(2) \backslash \Gamma$, $\Gamma = \text{SL}_2(\mathbb{Z})$. The summation over all γ gives

$$\begin{aligned} \langle \Delta(z), \Delta(2z) \rangle &= 2^{-6} [\Gamma : \Gamma_0(2)]^{-1} \sum_{\gamma} \langle \Delta(z) | \gamma, \Delta(z) | \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} \gamma \rangle \\ &= 2^{-6} 3^{-1} \left\langle \Delta(z), \text{Tr}^{(2)} \left(\Delta(z) | \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} \right) \right\rangle. \end{aligned} \quad (47)$$

The trace operator $\text{Tr}^{(N)} : \mathcal{M}_k(\Gamma_0(N)) \rightarrow \mathcal{M}_k(\text{SL}_2(\mathbb{Z}))$ is defined by the action $f \rightarrow \sum_{\gamma \in \Gamma_0(N) \backslash \text{SL}_2(\mathbb{Z})} f|_k \gamma$. We have (see [Ser73])

$$\text{Tr}^{(2)} \left(\Delta(z) | \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix} \right) = 2^{-5} T_2(\Delta) \quad (48)$$

where T_2 is the Hecke operator, therefore,

$$\begin{aligned} \langle \Delta(z), \Delta(2z) \rangle &= 2^{-6} 3^{-1} \langle \Delta(z), 2^{-5} T_2(\Delta(z)) \rangle \\ &= 2^{-11} 3^{-1} \langle \Delta(z), \tau(2) \Delta(z) \rangle \\ &= -\frac{1}{256} \langle \Delta(z), \Delta(z) \rangle. \end{aligned} \quad (49)$$

Substituting the last identity into (44), we obtain the final expression:

$$L(s, \Delta \otimes G_{2,2}) = \frac{3}{2} \frac{(4\pi)^{11}}{\Gamma(s)} \frac{(232 A_1(s) - A_2(s))}{256} \langle \Delta, \Delta \rangle. \quad (50)$$

5 Result for $L(s-9, \Delta) L(s-10, \Delta)$

Combining together (30) and (50) we obtain the expression for the product $L(s-9, \Delta) L(s-10, \Delta)$:

$$L(s-9, \Delta) L(s-10, \Delta) = \frac{3 \cdot 2^{13} \pi^{11} (232 A_1(s-9) - A_2(s-9))}{(1 + 3 \cdot 2^{13-s} + 2^{31-2s}) \Gamma(s-9)} \langle \Delta, \Delta \rangle. \quad (51)$$

Now we evaluate this result in the form $L(s-9, \Delta) L(s-10, \Delta) = R_\Delta(s) P_\Delta(s) \langle \Delta, \Delta \rangle$ for each $s \in \{12, \dots, 19\}$ computing the rational coefficient R_Δ and the corresponding power of π , see Table 2. The numerical value of the Petersson inner product

$$\langle \Delta, \Delta \rangle = 0.000001035362056 \quad (52)$$

is computed in the section 8.

Table 2: $L(s - 9, \Delta) L(s - 10, \Delta)$

s	R_Δ	P_Δ	$L(s - 9, \Delta) L(s - 10, \Delta)$
12	$\frac{32768}{225} = \frac{2^{15}}{3^2 \cdot 5^2}$	π^5	0.046143339818118
13	$\frac{4096}{81} = \frac{2^{12}}{3^4}$	π^7	0.158130732552033
14	$\frac{2048}{189} = \frac{2^{11}}{3^3 \cdot 7}$	π^9	0.334433094416363
15	$\frac{8192}{4725} = \frac{2^{13}}{3^3 \cdot 5^2 \cdot 7}$	π^{11}	0.528115574483468
16	$\frac{16384}{70875} = \frac{2^{14}}{3^4 \cdot 5^3 \cdot 7}$	π^{13}	0.694972239760782
17	$\frac{8192}{297675} = \frac{2^{13}}{3^5 \cdot 5^2 \cdot 7^2}$	π^{15}	0.816559651925946
18	$\frac{8192}{2679075} = \frac{2^{13}}{3^7 \cdot 5^2 \cdot 7^2}$	π^{17}	0.895457859377812
19	$\frac{65536}{200930625} = \frac{2^{16}}{3^8 \cdot 5^4 \cdot 7^2}$	π^{19}	0.942700248523234

6 Computation of $L(s, \Delta \otimes g_{20})$

We apply once again [Shi76, (2.4)] similarly as in section 4. The main difference is that the Petersson inner product is taken for both modular forms being cusp forms and for the full modular group $\mathrm{SL}_2(\mathbb{Z})$:

$$\begin{aligned}
 L(s, \Delta \otimes g_{20}) &= \frac{(4\pi)^s}{2\Gamma(s)} \int_{\Phi_1} \overline{g_{20}} \Delta E_{8,1}(z, s - 19) y^{s-1} dx dy \\
 &= \frac{(4\pi)^s}{2\Gamma(s)} \int_{\Phi_1} \overline{g_{20}} \Delta E_{8,1}(z, s - 19) y^{18} y^{s-19} dx dy \\
 &= \frac{(4\pi)^s}{2\Gamma(s)} \langle g_{20}, \Delta y^{s-19} E_{8,1}(z, s - 19) \rangle \\
 &= \frac{(4\pi)^s}{2\Gamma(s)} \langle g_{20}, \mathcal{H}ol(\Delta y^{s-19} E_{8,1}(z, s - 19)) \rangle \\
 &= \frac{(4\pi)^{19}}{2\Gamma(s)} \langle g_{20}, \mathcal{H}ol(\Delta (4\pi y)^{s-19} E_{8,1}(z, s - 19)) \rangle,
 \end{aligned} \tag{53}$$

In this case the critical values of s are $12, \dots, 19$. We verify that $s - 19 \leq 0$, $s - 19 + 8 = s - 11 > 0$, then the series $E_{8,1}(z, s - 19)$ is a nearly holomorphic modular form for all these $s \in \{12, \dots, 19\}$. We write the

Fourier expansion of $E_{8,1}(z, s-19)$ using [Pan03, Proposition 2.2]:

$$\begin{aligned}
& (4\pi y)^{s-19} E_{8,1}(z, s-19) \\
&= (4\pi y)^{s-19} \left[2\zeta(2s-30) \right. \\
&\quad + \frac{(-2\pi i)^{2s-30} (-1)^{s-19} \Gamma(2s-31)}{(4\pi y)^{2s-31} \Gamma(s-11) \Gamma(s-19)} 2\zeta(2s-31) \\
&\quad + \frac{2(-2\pi i)^{2s-30} (-1)^{s-19}}{\Gamma(s-11)} \times \\
&\quad \times \sum_{n=1}^{\infty} \sum_{d|n} d^{2s-31} W(4\pi n y, s-11, s-19) q^n \Big] \\
&= D'_0 (4\pi y)^{12-s} + D''_0 (4\pi y)^{s-19} \\
&\quad + \sum_{n=1}^{\infty} 2(2\pi)^{2s-30} \sum_{d|n} d^{2s-31} \frac{W(4\pi n y, s-11, s-19)}{\Gamma(s-11)} (4\pi y)^{s-11} q^n,
\end{aligned} \tag{54}$$

where

$$\begin{aligned}
D'_0 &= D'_0(s) = 2(2\pi)^{2s-30} \frac{\Gamma(2s-31) \zeta(2s-31)}{\Gamma(s-11) \Gamma(s-19)}, \\
D''_0 &= D''_0(s) = 2\zeta(2s-30).
\end{aligned} \tag{55}$$

Since the result of the holomorphic projection in this case belongs to the one-dimensional space spanned by g_{20} , we need to compute just the first Fourier coefficient $B_1(s)$ of $\mathcal{H}ol(\cdot) = \sum_{n=1}^{\infty} B_n(s) q^n$ in order to express it as a multiple of g_{20} . We compute it as the integral given by the Holomorphic Projection Lemma:

$$\begin{aligned}
B_1(s) &= \frac{1}{18!} \int_0^{+\infty} D''_0 (4\pi y)^{s-19} e^{-4\pi y} (4\pi y)^{18} d(4\pi y) \\
&\quad + \frac{1}{18!} \int_0^{+\infty} D'_0 (4\pi y)^{12-s} e^{-4\pi y} (4\pi y)^{18} d(4\pi y) \\
&= \frac{\Gamma(s)}{18!} D''_0 + \frac{\Gamma(31-s)}{18!} D'_0
\end{aligned} \tag{56}$$

The final expression is as following:

$$\begin{aligned}
L(s, \Delta \otimes g_{20}) &= B_1(s) \frac{(4\pi)^{19}}{2\Gamma(s)} \langle g_{20}, g_{20} \rangle \\
&= \left(\frac{\Gamma(s)}{18!} D''_0 + \frac{\Gamma(31-s)}{18!} D'_0 \right) \frac{(4\pi)^{19}}{2\Gamma(s)} \langle g_{20}, g_{20} \rangle \\
&= \frac{(4\pi)^{19}}{2 \cdot 18!} \left(D''_0 + \frac{\Gamma(31-s)}{\Gamma(s)} D'_0 \right) \langle g_{20}, g_{20} \rangle.
\end{aligned} \tag{57}$$

Now we evaluate this result for each $s \in \{12, \dots, 19\}$ in the form $L(s, \Delta \otimes g_{20}) = R_{g_{20}}(s) P_{g_{20}}(s) \langle g_{20}, g_{20} \rangle$ computing the rational coefficient $R_{g_{20}}$ and the corresponding power of π , see Table 3. The numerical value of the Petersson inner product

$$\langle g_{20}, g_{20} \rangle = 0.00000826554153165970 \tag{58}$$

is computed in the section 8.

Table 3: $L(s, \Delta \otimes g_{20})$

s	$R_{g_{20}}$	$P_{g_{20}}$	$L(s, \Delta \otimes g_{20})$
12	$\frac{524288}{2338875} = \frac{2^{19}}{3^5 \cdot 5^3 \cdot 7 \cdot 11}$	π^{13}	5.380003562880315
13	$\frac{2097152}{88409475} = \frac{2^{21}}{3^8 \cdot 5^2 \cdot 7^2 \cdot 11}$	π^{15}	5.618889612918517
14	$\frac{4194304}{2791213425} = \frac{2^{22}}{3^8 \cdot 5^2 \cdot 7 \cdot 11 \cdot 13 \cdot 17}$	π^{17}	3.513063561721911
15	$\frac{8388608}{97692469875} = \frac{2^{23}}{3^8 \cdot 5^3 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17}$	π^{19}	1.981288433718698
16	$\frac{8388608}{1465387048125} = \frac{2^{23}}{3^9 \cdot 5^4 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17}$	π^{21}	1.303635536350500
17	$\frac{2097152}{4396161144375} = \frac{2^{21}}{3^{10} \cdot 5^4 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17}$	π^{23}	1.072197252248449
18	$\frac{4194304}{92319384031875} = \frac{2^{22}}{3^{11} \cdot 5^4 \cdot 7^3 \cdot 11 \cdot 13 \cdot 17}$	π^{25}	1.007825020916877
19	$\frac{2097152}{461596920159375} = \frac{2^{21}}{3^{11} \cdot 5^5 \cdot 7^3 \cdot 11 \cdot 13 \cdot 17}$	π^{27}	0.994683426196918

7 The main identity

Combining (51) and (57) into the original expression (5) we get the following expression:

$$\begin{aligned}
 L(s, F_{12}, spin) &= L(s-9, \Delta) L(s-10, \Delta) L(s, \Delta \otimes g_{20}) \\
 &= \frac{3 \cdot 2^{13} \pi^{11}}{\Gamma(s-9)} \frac{(232 A_1(s-9) - A_2(s-9))}{(1 + 3 \cdot 2^{13-s} + 2^{31-2s})} \langle \Delta, \Delta \rangle \times \\
 &\quad \times \frac{(4\pi)^{19}}{2 \cdot 18!} \left(D_0''(s) + \frac{\Gamma(31-s)}{\Gamma(s)} D_0'(s) \right) \langle g_{20}, g_{20} \rangle = \quad (59) \\
 &= \frac{3 \cdot 2^{50} \pi^{30}}{18! \Gamma(s-9) (1 + 3 \cdot 2^{13-s} + 2^{31-2s})} \times \\
 &\quad \times \left(D_0''(s) + \frac{\Gamma(31-s)}{\Gamma(s)} D_0'(s) \right) \langle \Delta, \Delta \rangle \langle g_{20}, g_{20} \rangle,
 \end{aligned}$$

where $A_1(s)$ is given by (40), $A_2(s)$ is given by (41), D_0' and D_0'' are given by (55). For each critical value $s \in \{12, \dots, 19\}$ we evaluate this expression in the form $L(s, F_{12}, spin) = R(s) P(s) \langle \Delta, \Delta \rangle \langle g_{20}, g_{20} \rangle$ computing the rational coefficient R and the corresponding power of π , see Table 4.

Table 4: $L(s, F_{12}, spin)$

s	R	P	$L(s, F_{12}, spin)$
12	$\frac{17179869184}{526246875} = \frac{2^{34}}{3^7 \cdot 5^5 \cdot 7 \cdot 11}$	π^{18}	0.248251332624670
13	$\frac{8589934592}{7161167475} = \frac{2^{33}}{3^{12} \cdot 5^2 \cdot 7^2 \cdot 11}$	π^{22}	0.888519130619814
14	$\frac{8589934592}{527539337325} = \frac{2^{33}}{3^{11} \cdot 5^2 \cdot 7^2 \cdot 11 \cdot 13 \cdot 17}$	π^{26}	1.174884717828030
15	$\frac{68719476736}{461596920159375} = \frac{2^{36}}{3^{11} \cdot 5^3 \cdot 7^3 \cdot 11 \cdot 13 \cdot 17}$	π^{30}	1.046349279390801
16	$\frac{137438953472}{103859307035859375} = \frac{2^{37}}{3^{13} \cdot 5^7 \cdot 7^3 \cdot 11 \cdot 13 \cdot 17}$	π^{34}	0.905990508529256
17	$\frac{17179869184}{1308627268651828125} = \frac{2^{24}}{3^{15} \cdot 5^6 \cdot 7^4 \cdot 11 \cdot 13 \cdot 17}$	π^{38}	0.875513015091950
18	$\frac{34359738368}{247330553775195515625} = \frac{2^{35}}{3^{18} \cdot 5^6 \cdot 7^5 \cdot 11 \cdot 13 \cdot 17}$	π^{42}	0.902464835857626
19	$\frac{137438953472}{92748957665698318359375} = \frac{2^{37}}{3^{19} \cdot 5^9 \cdot 7^5 \cdot 11 \cdot 13 \cdot 17}$	π^{46}	0.937688313077777

8 Numerical computation of Petersson inner product

To compute numerically the Petersson inner product of Δ by itself and g_{20} by itself we use the classical result by Rankin [Ran52, Theorem 5]:

$$\langle f_k, f_k \rangle = \frac{(4\pi)^{1-k} (k-2)!}{\zeta(l)} \frac{\alpha_r}{\alpha_l + \alpha_r - \alpha_k} L(k-1, f_k) L(l, f_k), \quad (60)$$

where f_k is the cusp form of weight $k = \{12, 16, 18, 20, 22, 26\}$ of the form

$$f_k(z) = E_{k-12}(z) \Delta(z) \quad (61)$$

and

$$\begin{aligned} 4 &\leq r \leq k/2 - 2, \\ l &= k - r; \end{aligned} \quad (62)$$

E_k denotes Eisenstein series

$$\begin{aligned} E_k(z) &= \sum_{n=0}^{\infty} \alpha_k(n) q^n, \\ \alpha_k(0) &= 1, \\ \alpha_k &= \alpha_k(1) = -\frac{2k}{B_k}, \end{aligned} \quad (63)$$

B_k is a Bernoulli number.

For $f_k = \Delta$ we are able to use only one choice of critical value $l = 8$. To compute the numerical values $L(11, \Delta)$ and $L(8, \Delta)$ we used Dokchitser's L -functions Calculator [Dok]. In order to achieve the default precision (53 machine bits, which satisfies the functional equation to 1E-21), it is necessary to input 12 Fourier coefficients in this case. The obtained value is

$$\langle \Delta, \Delta \rangle = 0.000001035362056804320948209596804, \quad (64)$$

which coincides with the value given by Zagier in [Zag77, page 116] up to 11 digit (his method involves the direct summation of 250 first terms in L -series).

We used again Rankin's theorem to compute the Petersson inner product of g_{20} by itself. For the modular form of weight 20 there are three choices $l = 12, 14, 16$. For each choice of l we computed the special value of $L(l, g_{20})$ using Dokchitser's L -functions Calculator. It required to input 14 Fourier coefficients of g_{20} in order to achieve the default precision. The obtained values are

$$\begin{aligned} \langle g_{20}, g_{20} \rangle &= 0.000008265541531659702744699575969 \text{ for } l = 12, \\ \langle g_{20}, g_{20} \rangle &= 0.000008265541531659703390644766954 \text{ for } l = 14, \\ \langle g_{20}, g_{20} \rangle &= 0.000008265541531659703069998511729 \text{ for } l = 16. \end{aligned} \quad (65)$$

9 Numerical verification

The obtained values of $L(s, F_{12}, \text{spin})$ in section 7 can be numerically verified by using Dokchitser's L -functions Calculator by computing each term of the product in the righthand side of the identity (5). The computation of $L(s, \Delta)$ was already mentioned in the previous section.

To compute the values of $L(s, \Delta \otimes g_{20})$ for $s \in \{12, \dots, 19\}$ we have to determine the coefficients of this L -series first. Using the identity (16) for $f = \Delta = \sum \tau(n)n^{-s}$ and $g = g_{20} = \sum b(n)n^{-s} \in S_{20}$ we get

$$\begin{aligned} L(s, \Delta \otimes g_{20}) &= \sum_{n=1}^{\infty} A(n)n^{-s} \\ &= \sum_{d=1}^{\infty} d^{30-2s} \sum_{d_1=1}^{\infty} \tau(d_1)b(d_1)d_1^{-s} \\ &= \sum_{d, d_1 \geq 1} d^{30} \tau(d_1)b(d_1) (d^2 d_1)^{-s} \\ &= \sum_{n=1}^{\infty} \sum_{d: d^2 | n} d^{30} \tau\left(\frac{n}{d^2}\right) b\left(\frac{n}{d^2}\right) n^{-s}. \end{aligned} \quad (66)$$

Therefore we obtain

$$A(n) = \sum_{d: d^2 | n} d^{30} \tau\left(\frac{n}{d^2}\right) b\left(\frac{n}{d^2}\right). \quad (67)$$

The ComputeL program requires some functional equation parameters such as Γ -factors and the weight. These parameters can be deduced from

Table 5: Fourier coefficients of $L(s, \Delta \otimes g_{20})$

n	τ	g_{20}	A
1	1	1	1
2	-24	456	-10944
3	252	50652	12764304
4	-1472	-316352	1539411968
5	4830	-2377410	-11482890300
6	-6048	23097312	-139692542976
7	-16744	-16917544	283267356736
8	84480	-383331840	-44134904365056
9	-113643	1403363637	46408678295058
10	-115920	-1084098960	125668751443200
11	534612	-16212108	-8667187482096
12	-370944	-16023861504	19649522340790272
13	-577738	50421615062	-29130483042689756
14	401856	-7714400064	-3100077952118784
15	1217160	-120420571320	-146571102587851200

the Hodge structures (see [Sch90]) of Δ and g_{20} , namely

$$\begin{aligned}\Delta &\longrightarrow (0, 11) + (11, 0), \\ g_{20} &\longrightarrow (0, 19) + (19, 0).\end{aligned}\tag{68}$$

Therefore the Hodge structure of their tensor product (see [Yos01]) is

$$\Delta \otimes g_{20} \longrightarrow (0, 30) + (11, 19) + (19, 11) + (30, 0).\tag{69}$$

The Deligne's rule gives [Del79, page 329] in our case two Γ -factors: $\Gamma_{\mathbb{C}}(s)$ and $\Gamma_{\mathbb{C}}(s - 11)$. One can use the Gauss Duplication formula, which gives in our case four factors $\Gamma_{\mathbb{R}}(s) \Gamma_{\mathbb{R}}(s + 1) \Gamma_{\mathbb{R}}(s - 11) \Gamma_{\mathbb{R}}(s - 10)$. The weight that appears in the functional equation is the weight of the tensor product of two motives plus 1, which gives us $11 + 19 + 1 = 31$. We need about 150 coefficients of L -series to obtain the default precision (the functional equation is satisfied with 1E-27 precision). Coefficient of Δ and g_{20} are readily available in SAGE. First few coefficients $A(n)$ are given in Table 5.

Finally, we are able to compare the result in Table 4 and both of its parts in Tables 2, 3 with the direct numerical computation. These values and the absolute values of the difference with theoretical rational computation results are presented in Table 6 and Table 7.

Table 6: Numerical computation and comparison

s	$L(s-9, \Delta) L(s-10, \Delta)$	variation from Table 2	$L(s, \Delta \otimes g_{20})$	variation from Table 3
12	0.046143339853964	3.58E-11	5.38000356288032	4.95E-15
13	0.158130732674877	1.23E-10	5.61888961291852	2.39E-15
14	0.334433094676168	2.60E-10	3.51306356172191	1.44E-15
15	0.528115574893734	4.10E-10	1.98128843371870	1.36E-15
16	0.694972240300672	5.40E-10	1.30363553635050	7.99E-16
17	0.816559652560290	6.34E-10	1.07219725224845	7.40E-16
18	0.895457860073449	6.96E-10	1.00782502091688	2.75E-15
19	0.942700249255570	7.32E-10	0.99468342619692	4.22E-16

Table 7: Numerical computation and comparison (final)

s	$L(s, F_{12}, spin)$	variation from Table 4
12	0.24825133281752	1.98E-10
13	0.88851913131006	6.90E-10
14	1.17488471874074	9.13E-10
15	1.04634928020366	8.13E-10
16	0.90599050923308	7.04E-10
17	0.87551301577209	6.80E-10
18	0.90246483655871	7.01E-10
19	0.93768831380622	7.28E-10

References

- [CP04] Michel Courtieu and Alexei Panchishkin. *Non-Archimedean L -functions and arithmetical Siegel modular forms*, volume 1471 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, second edition, 2004.
- [Del79] Pierre Deligne. Valeurs de fonctions L et périodes d'intégrales. In *Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2*, Proc. Sympos. Pure Math., XXXIII, pages 313–346. Amer. Math. Soc., Providence, R.I., 1979. With an appendix by N. Koblitz and A. Ogus.
- [Dok] Tim Dokchitser. *ComputeL – Computing special values of L -functions*. <http://www.maths.dur.ac.uk/~dma0td/computel/>. v1.3.
- [Dok04] Tim Dokchitser. Computing special values of motivic L -functions. *Experiment. Math.*, 13(2):137–149, 2004.
- [GZ86] Benedict H. Gross and Don B. Zagier. Heegner points and derivatives of L -series. *Invent. Math.*, 84(2):225–320, 1986.
- [Hei08] Bernhard Heim. Miyawaki’s F_{12} spinor L -function conjecture. 2008. arXiv:0712.1286v1 [math.NT].
- [Ike06] Tamotsu Ikeda. Pullback of the lifting of elliptic cusp forms and Miyawaki’s conjecture. *Duke Math. J.*, 131(3):469–497, 2006.
- [Miy92] Isao Miyawaki. Numerical examples of Siegel cusp forms of degree 3 and their zeta-functions. *Mem. Fac. Sci. Kyushu Univ. Ser. A*, 46(2):307–339, 1992.
- [Miy06] Toshitsune Miyake. *Modular forms*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, english edition, 2006. Translated from the 1976 Japanese original by Yoshitaka Maeda.
- [Pan03] A. A. Panchishkin. Two variable p -adic L functions attached to eigenfamilies of positive slope. *Invent. Math.*, 154(3):551–615, 2003.
- [Ran52] R. A. Rankin. The scalar product of modular forms. *Proc. London Math. Soc. (3)*, 2:198–217, 1952.
- [SAG] *SAGE Mathematical Software*. <http://www.sagemath.org>. Ver. 2.11.
- [Sch90] A.J. Scholl. Motives for modular forms. *Invent. Math.*, 100(2):419–430, 1990.
- [Ser73] Jean-Pierre Serre. Formes modulaires et fonctions zêta p -adiques. In *Modular functions of one variable, III (Proc. Internat. Summer School, Univ. Antwerp, 1972)*, pages 191–268. Lecture Notes in Math., Vol. 350. Springer, Berlin, 1973.
- [Shi76] Goro Shimura. The special values of the zeta functions associated with cusp forms. *Comm. Pure Appl. Math.*, 29(6):783–804, 1976.
- [Shi07] Goro Shimura. *Elementary Dirichlet series and modular forms*. Springer Monographs in Mathematics. Springer, New York, 2007.

- [Stu80] Jacob Sturm. Projections of C^∞ automorphic forms. *Bull. Amer. Math. Soc. (N.S.)*, 2(3):435–439, 1980.
- [Yos01] Hiroyuki Yoshida. Motives and Siegel modular forms. *Amer. J. Math.*, 123(6):1171–1197, 2001.
- [Zag77] Don Zagier. Modular forms whose Fourier coefficients involve zeta-functions of quadratic fields. In *Modular functions of one variable, VI (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976)*, pages 105–169. Lecture Notes in Math., Vol. 627. Springer, Berlin, 1977.
- [Žar74] N. A. Žarkovskaja. The Siegel operator and Hecke operators. *Funkcional. Anal. i Priložen.*, 8(2):30–38, 1974. in Russian.